

Medical Policy

MRI of the Spine

Policy Number: 1027

Policy History

Approve Date:	11/14/2014	Effective Date:	11/14/2014
---------------	------------	-----------------	------------

Preauthorization

All Plans	Benefit plans vary in coverage and some plans may not provide coverage for certain service(s) listed in this policy. Decisions for authorization are subject to all terms and conditions of the applicable benefit plan, including specific exclusions and limitations as well as applicable state and/or federal laws. Please review the benefit plan descriptions for details.
-----------	--

Policy

Indications of Coverage

- I. WEA Trust considers MRI of the cervical and thoracic and lumbar spine, medically necessary for the following indications when the corresponding criteria are met:
- II. Thoracic, or cervical pain that is localized or radicular in nature as indicated by ALL of the following:
 - A. Pain presents with objective findings of motor or reflex changes in the specific nerve root distribution
 - B. Member has tried and failed to improve after six weeks of conservative treatment for the current episode of pain including all of the following:
 - i. Analgesics, NSAIDs, or oral corticosteroids
 - ii. Physical therapy
 - iii. Modification of activity
 - C. Member is being considered for invasive treatment (e.g. epidural steroids or surgery)
 - D. Pain significantly interferes with member's daily function
- III. Lumbar pain that is localized or radicular in nature as indicated by ALL of the following:
 - A. Pain presents with objective findings of motor or reflex changes in the specific nerve root distribution
 - B. Member has tried and failed to improve after six weeks of conservative treatment for the current episode of pain including all of the following:
 - i. Analgesics, NSAIDs, or oral corticosteroids
 - ii. Physical therapy or participation in the Kiio program
 - iii. Modification of activity
 - C. Member is being considered for invasive treatment (e.g. epidural steroids or surgery)
 - D. Pain significantly interferes with member's daily function
- IV. Severe pain (level 8-10/10 on pain scale) that meets both of the following:
 - A. Unresponsive to oral medications including NSAIDs, a dosepak, muscle relaxants, and narcotics
 - B. Significantly limits ability to perform daily activities and interferes with sleep

- V. Suspected spinal stenosis as indicated by ALL of the following:
 - A. Neurogenic claudication with progressive or disabling symptoms indicated by one or more of the following:
 - i. Lumbar spine
 - 1. Progressive or disabling symptoms of lumbar spine stenosis, as indicated by one or more of the following
 - a. Back, buttock, or leg pain worsening with prolonged standing and lumbar extension
 - b. Leg weakness
 - c. Wide-based gait or abnormal Romberg test
 - ii. Cervical and Thoracic spine
 - 1. Progressive or disabling symptoms of cervical or thoracic spine stenosis, as indicated by one or more of the following.
 - a. Hyperactive reflexes
 - b. Muscle weakness
 - c. Sensory loss
 - d. Spasticity
 - B. Member is being considered for invasive treatment
- VI. Known or suspected spondylolysis by radiologic evidence (e.g. by plain x-ray, bone scan, CT scan), with or without spondylolisthesis, and one or more of the following:
 - A. Focal neurogenic findings
 - B. Significant pain
 - C. Urinary retention or incontinence
- VII. Suspected spinal cord compression, myelopathy, or cauda equine syndrome as indicated by one or more of the following:
 - A. Babinski sign positive (cervical spine)
 - B. Fecal and/or urinary incontinence
 - C. Gait abnormality
 - D. Hoffman sign positive (cervical spine)
 - E. Hyperreflexia or clonus
 - F. Significant or progressive sensory or motor deficits
 - G. Saddle anesthesia
 - H. Spasticity
- VIII. Known or suspected fracture/dislocation of spine or spinal cord injury due to trauma as indicated by one or more of the following:
 - A. Myelopathy
 - B. Neurologic findings corresponding to vertebral level of injury seen on x-ray or CT scan
 - C. Progressive neurological deficit
 - D. Radiculopathy
 - E. Neurological symptoms with traumatic mechanism of injury such as fall from a great height, motor vehicle accident, etc.
 - F. Significant head injury (cervical spine)
 - G. Clinical suspicion of acute spinal injury as indicated by one or more of the following
 - i. Negative or indeterminate spine x-ray
 - ii. Symptoms such as midline cervical tenderness or paresthesia of extremities
 - H. Suspected epidural hematoma (cervical spine)
 - I. Suspected traumatic disk herniation
 - J. X-ray shows abnormal spine motion or burst fracture

- IX. Known or suspected ankylosing spondylitis or other seronegative spondyloarthritis as indicated by ALL of the following:
 - A. Back pain improving with exercise
 - B. Back pain of insidious onset that has persisted for three months or longer
 - C. Early-morning stiffness lasting longer than 30 minutes
 - D. Patient evaluated by rheumatologist
 - E. Plain x-ray results are nondiagnostic

- X. Cancer or neoplasm evaluation/staging as indicated by one or more of the following:
 - A. Localized back or neck pain and one or more of the following
 - i. Positive bone scan
 - ii. Night pain
 - iii. Persistent back pain in members older than 50 years
 - iv. Personal history or concurrent diagnosis of malignancy
 - v. Rapidly progressing weakness
 - vi. Unexplained weight loss
 - B. Myelopathy signs or symptoms (e.g. motor weakness, bowel/bladder dysfunction)
 - C. Post-treatment monitoring of spinal metastasis

- XI. Stereotactic spine radiotherapy treatment planning

- XII. Known or suspected infection (e.g. vertebral osteomyelitis, epidural abscess) as indicated by ALL of the following:
 - A. Localized midline back pain
 - B. Risk factors for spinal infection such as
 - i. Bone scan or plain x-ray suggest infection
 - ii. Elevated sedimentation rate
 - iii. Fever
 - iv. Recent invasive procedure
 - v. Immunosuppression
 - vi. Recent spinal surgery
 - vii. Tuberculosis, concurrent or suspected

- XIII. Suspected inflammatory or demyelinating process as indicated by one or more of the following.
 - A. Multiple sclerosis (MS) as indicated by one or more the following:
 - i. Ascending numbness or tingling
 - ii. Brown-Sequard syndrome
 - iii. Conditions similar to MS (Sjorgren syndrome, lupus, etc.) cannot be ruled out
 - iv. Brain MRI is negative for MS
 - v. Signs or symptoms of myelopathy or myelitis
 - B. Suspected transverse myelitis as indicated by one or more of the following:
 - i. Bilateral signs/symptoms of spinal cord of appropriate level of spinal cord
 - ii. Clearly defined sensory level
 - iii. Sudden onset of sensory, motor, and autonomic dysfunction attributable to appropriate level of spinal cord

- XIV. Suspected syringomyelia as indicated by:
 - A. Lumbar and thoracic spine weakness, muscle wasting, or sensory loss in appropriate spine dermatomes
 - B. Cervical spine upper limb sensory loss in cape-like distribution or weakness, muscle wasting, or sensory loss beginning in intrinsic muscles of hand

- XV. Scoliosis as indicated by one or more of the following:
 - A. Congenital scoliosis

- B. Early onset before five years of age
- C. Neurofibromatosis
- D. Pre-surgical planning for adolescent members to assess possible neural axis malformation as indicated by one or more of the following:
 - i. Abnormal neurological finding during clinical exam
 - ii. Age at first visit 10 years or younger
 - iii. Kyphosis at curve apex
 - iv. Left-sided thoracic curvature
 - v. Male
 - vi. Moderate to severe pain
 - vii. Short segment curve (<6 vertebral segments)
 - viii. Thoracic kyphosis is ≥ 30 degrees

XVI. Known or suspected congenital spinal conditions including:

- A. Anterior sacral meningocele
- B. Caudal regressing syndrome
- C. Chiari malformation (cervical spine)
- D. Diastematomyelia
- E. Dorsal dermal sinus
- F. Hydromyelia
- G. Intradural lipoma
- H. Klippel-Feil syndrome (cervical spine)
- I. Myelocele
- J. Myelomeningocele
- K. Spinal dysraphism
- L. Split notochord syndrome
- M. Tight filum terminale

XVII. Known or suspected tuberculosis of spine as indicated by one or more of the following:

- A. Diagnostic evaluation of suspected spinal tuberculosis
- B. Preparation of surgical management

XVII. Suspected tethered cord (lumbar spine) as indicated by one or more of the following:

- A. Anorectal malformation
- B. Cutaneous manifestation of occult spina bifida
- C. Gait abnormality or difficulty
- D. Lack of bladder control
- E. Abnormal urodynamic tests

XVIII. Torticollis (cervical spine) as indicated by ALL of the following:

- A. Member younger than 18 years
- B. Initial CT scan normal or inconclusive
- C. Nontraumatic etiology

XIX. Rheumatoid arthritis in cervical spine with suspected instability or cord compression as indicated by one or more of the following:

- A. Atlantoaxial subluxation or impaction of plain x-ray
- B. Myelopathy or progressive neurological deficit
- C. Subluxation of C1 and C2

XX. Repeat evaluation of specific area or structure with the same imaging modality as indicated by one or more of the following:

- A. New or worsening symptoms

- B. Interval reassessment may impact treatment plan
- C. Need for reimaging prior to or after invasive procedure

XXI. WEA Trust considers MRI of the spine not medically necessary for all other indications because of lack of clinical value. Clinical guidelines from several health agencies (Agency for Healthcare Policy and Research, American College of Physicians, and North American Spine Society) recommend against imaging studies for acute low back pain.

Background

When used appropriately, MRI (magnetic resonance imaging) is an effective diagnostic tool. Because bulging disks and other spinal abnormalities have been found in over half of all adults who are otherwise asymptomatic, it is important for clinicians to order MRIs judiciously and interpret results in the context of clinical findings and patient symptoms in order to avoid unnecessary treatment of conditions that may not be causing symptoms. This is especially relevant with the presence of low back pain without radiculopathy, severe progressive neurological deficits, trauma, or clinical suspicion of a severe underlying condition (cancer, infection, etc.). MRI is not necessary in these cases unless the back pain is severe (hospitalization is required) or when symptoms are progressive despite conservative treatment.

The American College of Physicians (2012) and North American Spine Society recommend against obtaining imaging studies in patients with non-specific back pain within the first six weeks of the onset of symptoms, unless red flags are present. Red flags include severe or progressive neurological deficits or when serious underlying conditions are suspected. Imaging of the low back before six weeks does not improve patient outcomes. Conservative treatment for spine-related pain is recommended for the first six weeks. Standard conservative care includes, but is not limited to, physical therapy, NSAIDs, narcotic or nonnarcotic analgesics, oral corticosteroid course, spinal manipulation, activity modification, rest, heat or ice, and cervical collars or braces. If conservative treatment fails to improve symptoms, MRI may be indicated to help guide future, possibly more invasive treatment.

References

The above policy is based on the following references:

1. Abdel-Wanis ME, Solyman MT, Hasan NM. Sensitivity, Specificity and Accuracy of Magnetic Resonance Imaging for Differentiating Vertebral Compression Fractures Caused by Malignancy, Osteoporosis, and Infections. *Journal of Orthopaedic Surgery (Hong Kong)* 2011;19(2):145-50.
2. Ackland HM, et al. Cervical Spine Magnetic Resonance Imaging in Alert, Neurologically Intact Trauma Patients with Persistent Midline Tenderness and Negative Computed Tomography Results. *Annals of Emergency Medicine* 2011;58(6):521-30. DOI: 10.1016/j.annemergmed.2011.06.008.
3. ACR-ASNR-SCBT-MR Practice Guideline for the Performance of Magnetic Resonance Imaging (MRI) of the Adult Spine. [Internet] American College of Radiology. Amended 2014 Accessed at: <http://www.acr.org/>. [created 2001; accessed 2014 Sept 23]
4. Alper G, Petropoulou KA, Fitz CR, Kim Y. Idiopathic Acute Transverse Myelitis in Children: an Analysis and Discussion of MRI Findings. *Multiple Sclerosis* 2011;17(1):74-80. DOI: 10.1177/1352458510381393.
5. Amrami KK. Imaging of the Seronegative Spondyloarthropathies. *Radiologic Clinics of North America* 2012;50(4):841-54. DOI: 10.1016/j.rcl.2012.04.010.
6. Andronikou S, et al. Musculoskeletal Tuberculosis - Imaging Using Low-End and Advanced Modalities for Developing and Developed Countries. *Acta Radiologica* 2011;52(4):430-41. DOI: 10.1258/ar.2011.100444.
7. Beattie PF. Diffusion-Weighted Magnetic Resonance Imaging of the Musculoskeletal System: an Emerging Technology with Potential to Impact Clinical Decision Making. *Journal of Orthopaedic and Sports Physical Therapy* 2011;41(11):887-95. DOI: 10.2519/jospt.2011.3744.
8. Boden BP, Jarvis CG. Spinal Injuries in Sports. *Neurologic Clinics* 2008;26(1):63-78; viii. DOI: 10.1016/j.ncl.2007.12.005.
9. Cardoso M, Keating RF. Neurosurgical Management of Spinal Dysraphism and Neurogenic Scoliosis. *Spine* 2009;34(17):1775-82. DOI: 10.1097/BRS.0b013e3181b07914.
10. Ceccarelli A, Bakshi R, Neema M. MRI in Multiple Sclerosis: a Review of the Current Literature. *Current Opinion in Neurology* 2012;25(4):402-9. DOI: 10.1097/WCO.0b013e328354f63f.
11. Chandra J, et al. MRI in Acute and Subacute Post-Traumatic Spinal Cord Injury: Pictorial Review. *Spinal Cord* 2012;50(1):2-7. DOI: 10.1038/sc.2011.107.

12. Chiapparini L, Saletti V, Solero CL, Bruzzone MG, Valentini LG. Neuroradiological Diagnosis of Chiari Malformations. *Neurological Sciences* 2011;32 Suppl 3:283-6. DOI: 10.1007/s10072-011-0695-0.
13. Chou D, et al. Degenerative Magnetic Resonance Imaging Changes in Patients with Chronic Low Back Pain: A Systematic Review. *Spine* 2011;36(21 Suppl):S43-53. DOI: 10.1097/BRS.0b013e31822ef700.
14. Chou R, Deyo RA, Jarvik JG. Appropriate Use of Lumbar Imaging for Evaluation of Low Back Pain. *Radiologic Clinics of North America* 2012;50(4):569-85. DOI: 10.1016/j.rcl.2012.04.005.
15. Chronic Neck Pain. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2013 Oct Accessed at: <http://www.acr.org>. [created 1998; accessed 2014 Sept 23]
16. Cox BW, et al. International Spine Radiosurgery Consortium Consensus Guidelines for Target Volume Definition in Spinal Stereotactic Radiosurgery. *International Journal of Radiation Oncology, Biology and Physics* 2012;83(5):e597-605. DOI: 10.1016/j.ijrobp.2012.03.009.
17. Crocker M, Jones TL, Rich P, Bell BA, Papadopoulos MC. The Clinical Value of Early Postoperative MRI After Lumbar Spine Surgery. *British Journal of Neurosurgery* 2010;24(1):46-50. DOI: 10.3109/02688690903374067.
18. Currie S, Galea-Soler S, Barron D, Chandramohan M, Groves C. MRI Characteristics of Tuberculous Spondylitis. *Clinical Radiology* 2011;66(8):778-87. DOI: 10.1016/j.crad.2011.02.016.
19. Cyteval C, Bourdon A. Imaging Orthopedic Implant Infections. *Diagnostic and Interventional Imaging* 2012;93(6):547-57. DOI: 10.1016/j.diii.2012.03.004.
20. Dahele M, et al. Imaging for Stereotactic Spine Radiotherapy: Clinical Considerations. *International Journal of Radiation Oncology, Biology and Physics* 2011;81(2):321-30. DOI: 10.1016/j.ijrobp.2011.04.039.
21. Del Grande F, Maus TP, Carrino JA. Imaging the Intervertebral Disk: Age-Related Changes, Herniations, and Radicular Pain. *Radiologic Clinics of North America* 2012;50(4):629-49. DOI: 10.1016/j.rcl.2012.04.014.
22. Demertzis JL, Rubin DA. MR Imaging Assessment of Inflammatory, Crystalline-Induced, and Infectious Arthritides. *Magnetic Resonance Imaging Clinics of North America* 2011;19(2):339-63. DOI: 10.1016/j.mric.2011.02.004.
23. DeSanto J, Ross JS. Spine Infection/Inflammation. *Radiologic Clinics of North America* 2011;49(1):105-27. DOI: 10.1016/j.rcl.2010.07.018.
24. Diab M, et al. Use and Outcome of MRI in the Surgical Treatment of Adolescent Idiopathic Scoliosis. *Spine* 2011;36(8):667-71. DOI: 10.1097/BRS.0b013e3181da218c.
25. Diaz JJ, et al. Practice Management Guidelines for the Screening of Thoracolumbar Spine Fracture. *Journal of Trauma* 2007;63(3):709-18. DOI: 10.1097/TA.0b013e318142d2db.
26. Diehn FE. Imaging of Spine Infection. *Radiologic Clinics of North America* 2012;50(4):777-98. DOI: 10.1016/j.rcl.2012.04.001.
27. Dundamadappa SK, Cauley KA. MR Imaging of Acute Cervical Spinal Ligamentous and Soft Tissue Trauma. *Emergency Radiology* 2012;19(4):277-86. DOI: 10.1007/s10140-012-1033-4.
28. Dunn R, Zondagh I, Candy S. Spinal Tuberculosis: Magnetic Resonance Imaging and Neurological Impairment. *Spine* 2011;36(6):469-73.
29. Easter JS, Barkin R, Rosen CL, Ban K. Cervical Spine Injuries in Children, Part II: Management and Special Considerations. *Journal of Emergency Medicine* 2011;41(3):252-6. DOI: 10.1016/j.jemermed.2010.03.018.
30. Elliott JM, Flynn TW, Al-Najjar A, Press J, Nguyen B, Noteboom JT. The Pearls and Pitfalls of Magnetic Resonance Imaging for the Spine. *Journal of Orthopaedic and Sports Physical Therapy* 2011;41(11):848-60. DOI: 10.2519/jospt.2011.3636.
31. Emch TM, Modic MT. Imaging of Lumbar Degenerative Disk Disease: History and Current State. *Skeletal Radiology* 2011;40(9):1175-89. DOI: 10.1007/s00256-011-1163-x.
32. Filippi M, et al. Association Between Pathological and MRI Findings in Multiple Sclerosis. *Lancet Neurology* 2012;11(4):349-60. DOI: 10.1016/S1474-4422(12)70003-0.
33. Filippi M, et al. Magnetic Resonance Techniques in Multiple Sclerosis: The Present and the Future. *Archives of Neurology* 2011;68(12):1514-20. DOI: 10.1001/archneurol.2011.914.
34. Filippi M, Rocca MA. MR Imaging of Multiple Sclerosis. *Radiology* 2011;259(3):659-81. DOI: 10.1148/radiol.11101362.
35. Fucs PM, Meves R, Yamada HH. Spinal Infections in Children: A Review. *International Orthopaedics* 2012;36(2):387-95. DOI: 10.1007/s00264-011-1388-2.

36. Furlan JC, Bracken MB, Fehlings MG. Is Age a Key Determinant of Mortality and Neurological Outcome After Acute Traumatic Spinal Cord Injury? *Neurobiology of Aging* 2010;31(3):434-46. DOI: 10.1016/j.neurobiolaging.2008.05.003.
37. Furlan JC, Noonan V, Cadotte DW, Fehlings MG. Timing of Decompressive Surgery of Spinal Cord After Traumatic Spinal Cord Injury: An Evidence-Based Examination of Pre-Clinical and Clinical Studies. *Journal of Neurotrauma* 2011;28(8):1371-99. DOI: 10.1089/neu.2009.1147.
38. George TM, Higginbotham NH. Defining the Signs and Symptoms of Chiari Malformation Type I With and Without Syringomyelia. *Neurological Research* 2011;33(3):240-6. DOI: 10.1179/016164111X12962202723760.
39. Goertz M, et al. Adult Acute and Subacute Low Back Pain. Health Care Guideline [Internet] Institute for Clinical Systems Improvement. 15th ed; 2012 Nov Accessed at: <http://www.icsi.org/>. [accessed 2014 Sept 23]
40. Graber JJ, Nolan CP. Myelopathies in Patients with Cancer. *Archives of Neurology* 2010;67(3):298-304. DOI: 10.1001/archneurol.2010.20.
41. Haque S, Bilal Shafi BB, Kaleem M. Imaging of Torticollis in Children. *Radiographics* 2012;32(2):557-71. DOI: 10.1148/rg.322105143.
42. Hermann KG, et al. Descriptions of Spinal MRI Lesions and Definition of a Positive MRI of the Spine in Axial Spondyloarthritis: A Consensual Approach by the ASAS/OMERACT MRI Study Group. *Annals of the Rheumatic Diseases* 2012;71(8):1278-88. DOI: 10.1136/ard.2011.150680.
43. Hodel J, Outteryck O, Jissendi P, Zins M, Leclerc X, Pruvo JP. MRI of Myelitis. *JBR-BTR : Organe de la Societe Royale Belge de Radiologie (SRBR)* 2012;95(4):270-6.
44. Jacob A, Weinshenker BG. An Approach to the Diagnosis of Acute Transverse Myelitis. *Seminars in Neurology* 2008;28(1):105-20. DOI: 10.1055/s-2007-1019132.
45. Johnston CE. Preoperative Medical and Surgical Planning for Early Onset Scoliosis. *Spine* 2010;35(25):2239-44. DOI: 10.1097/BRS.0b013e3181fd5853.
46. Jones TM, Anderson PA, Noonan KJ. Pediatric Cervical Spine Trauma. *Journal of the American Academy of Orthopedic Surgeons* 2011;19(10):600-11.
47. Kang Y, et al. New MRI Grading System for the Cervical Canal Stenosis. *American Journal of Roentgenology* 2011;197(1):W134-40. DOI: 10.2214/AJR.10.5560.
48. Karrar EE, Mansour N, Bhansali A. Cranial and Spinal Trauma: Current Concepts. *Disease-a-Month* 2011;57(10):543-57. DOI: 10.1016/j.disamonth.2011.08.025.
49. Katz JN, Harris MB. Clinical Practice. Lumbar Spinal Stenosis. *New England Journal of Medicine* 2008;358(8):818-25. DOI: 10.1056/NEJMcp0708097.
50. Khalil JG, Nassr A, Maus TP. Physiologic Imaging of the Spine. *Radiologic Clinics of North America* 2012;50(4):599-611. DOI: 10.1016/j.rcl.2012.04.004.
51. Kilsdonk ID, de Graaf WL, Barkhof F, Wattjes MP. Inflammation High-Field Magnetic Resonance Imaging. *Neuroimaging Clinics of North America* 2012;22(2):135-57, ix. DOI: 10.1016/j.nic.2012.02.010.
52. Kim H, et al. Scoliosis Imaging: What Radiologists Should Know. *Radiographics* 2010;30(7):1823-42. DOI: 10.1148/rg.307105061.
53. Kim HJ, Green DW. Spondylolysis in the Adolescent Athlete. *Current Opinion in Pediatrics* 2011;23(1):68-72. DOI: 10.1097/MOP.0b013e32834255c2.
54. Kim NH, Cho KT, Seo HS. Myelopathy Due to Intracranial Dural Arteriovenous Fistula: A Potential Diagnostic Pitfall. Case report. *Journal of Neurosurgery* 2011;114(3):830-3. DOI: 10.3171/2010.5.JNS10128.
55. Kim SH, Smith SE, Mulligan ME. Hematopoietic Tumors and Metastases Involving Bone. *Radiologic Clinics of North America* 2011;49(6):1163-83, vi. DOI: 10.1016/j.rcl.2011.07.004.
56. Knoeller SM, Uhl M, Gahr N, Adler CP, Herget GW. Differential Diagnosis of Primary Malignant Bone Tumors in the Spine and Sacrum. *The Radiological and Clinical Spectrum: Minireview. Neoplasma* 2008;55(1):16-22.
57. Korteweg T, et al. A Search for New MRI Criteria for Dissemination in Space in Subjects with a Clinically Isolated Syndrome. *European Radiology* 2009;19(9):2244-8. DOI: 10.1007/s00330-009-1402-x.
58. Kotsenas AL. Imaging of Posterior Element Axial Pain Generators: Facet Joints, Pedicles, Spinous Processes, Sacroiliac Joints, and Transitional Segments. *Radiologic Clinics of North America* 2012;50(4):705-30. DOI: 10.1016/j.rcl.2012.04.008.
59. Kreiner DS, et al. Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care [Internet] North American Spine Society. 2011 Accessed at: <http://www.spine.org/>.

60. Law M. Neurological Complications. *Cancer Imaging* 2009;9 Spec No A:S71-4. DOI: 10.1102/1470-7330.2009.9030.
61. Leone A, Cianfoni A, Cerase A, Magarelli N, Bonomo L. Lumbar Spondylolysis: A Review. *Skeletal Radiology* 2010;30(4):209-23. DOI: 10.1007/s00256-010-0942-0.
62. Leone A, et al. Imaging of Spondylodiscitis. *European Review for Medical and Pharmacological Sciences* 2012;16 Suppl 2:8-19.
63. Lew SM, Kothbauer KF. Tethered Cord Syndrome: An Updated Review. *Pediatric Neurosurgery* 2007;43(3):236-48. DOI: 10.1159/000098836.
64. Lindberg PG, Feydy A, Sanchez K, Rannou F, Maier MA. Measures of Spinal Canal Stenosis and Relationship to Spinal Cord Structure in Patients with Cervical Spondylosis. *Journal of Neuroradiology* 2012;39(4):236-42. DOI: 10.1016/j.neurad.2011.09.004.
65. Looby S, Flanders A. Spine Trauma. *Radiologic Clinics of North America* 2011;49(1):129-63. DOI: 10.1016/j.rcl.2010.07.019.
66. Lovblad KO, et al. MR Imaging in Multiple Sclerosis: Review and Recommendations for Current Practice. *AJNR. American Journal of Neuroradiology* 2009;31(6):983-9. DOI: 10.3174/ajnr.A1906.
67. Low Back Pain. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2011 Accessed at: <http://www.acr.org/>. [created 1996; accessed 2014 Sept 23]
68. Maksymowych WP, Weber U. Diagnostic Utility of MRI in Early Spondyloarthritis. *Current Rheumatology Reports* 2011;13(5):402-8. DOI: 10.1007/s11926-011-0190-x.
69. Malfair D, et al. Radiographic Evaluation of Scoliosis: Review. *American Journal of Roentgenology* 2010;194(3 Suppl):S8-22. DOI: 10.2214/AJR.07.7145.
70. Maus T. Imaging the Back Pain Patient. *Physical Medicine and Rehabilitation Clinics of North America* 2010;21(4):725-66. DOI: 10.1016/j.pmr.2010.07.004.
71. Maus TP, Aprill CN. Lumbar Diskogenic Pain, Provocation Diskography, and Imaging Correlates. *Radiologic Clinics of North America* 2012;50(4):681-704. DOI: 10.1016/j.rcl.2012.04.013.
72. Maus TP. Imaging of Spinal Stenosis: Neurogenic Intermittent Claudication and Cervical Spondylotic Myelopathy. *Radiologic Clinics of North America* 2012;50(4):651-79. DOI: 10.1016/j.rcl.2012.04.007.
73. McNee P, et al. Predictors of Long-Term Pain and Disability in Patients with Low Back Pain Investigated by Magnetic Resonance Imaging: A Longitudinal Study. *BMC Musculoskeletal Disorders* 2011;12:234. DOI: 10.1186/1471-2474-12-234.
74. Metastatic Bone Disease. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2012 Accessed at: <http://www.acr.org/>. [created 1995; accessed 2014 Sept 23]
75. Montalban X, et al. MRI Criteria for MS in Patients with Clinically Isolated Syndromes. *Neurology* 2010;74(5):427-34. DOI: 10.1212/WNL.0b013e3181cec45c.
76. Murthy NS. Imaging of Stress Fractures of the Spine. *Radiologic Clinics of North America* 2012;50(4):799-821. DOI: 10.1016/j.rcl.2012.04.009.
77. Myelopathy. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2011 Accessed at: <http://www.acr.org/>. [created 1996; accessed 2014 Sept 23]
78. Naganawa T, Miyamoto K, Ogura H, Suzuki N, Shimizu K. Comparison of Magnetic Resonance Imaging and Computed Tomogram-Myelography for Evaluation of Cross Sections of Cervical Spinal Morphology. *Spine* 2011;36(1):50-6. DOI: 10.1097/BRS.0b013e3181cb469c.
79. Ogura H, Miyamoto K, Fukuta S, Naganawa T, Shimizu K. Comparison of Magnetic Resonance Imaging and Computed Tomography-Myelography for Quantitative Evaluation of Lumbar Intracanalicular Cross-Section. *Yonsei Medical Journal* 2011;52(1):137-44. DOI: 10.3349/ymj.2011.52.1.137.
80. Pizones J, et al. Impact of Magnetic Resonance Imaging on Decision Making for Thoracolumbar Traumatic Fracture Diagnosis and Treatment. *European Spine Journal* 2011;20 Suppl 3:390-6. DOI: 10.1007/s00586-011-1913-4.
81. Polman CH, et al. Diagnostic Criteria for Multiple Sclerosis: 2010 Revisions to the McDonald Criteria. *Annals of Neurology* 2011;69(2):292-302. DOI: 10.1002/ana.22366.
82. Rihn JA, Harrod C, Albert TJ. Revision Cervical Spine Surgery. *Orthopedic Clinics of North America* 2012;43(1):123-36, ix-x. DOI: 10.1016/j.ocl.2011.09.001.
83. Roudsari B, Jarvik JG. Lumbar Spine MRI for Low Back Pain: Indications and Yield. *American Journal of Roentgenology* 2010;195(3):550-9. DOI: 10.2214/AJR.10.4367.

84. Roy AK, Slimack NP, Ganju A. Idiopathic Syringomyelia: Retrospective Case Series, Comprehensive Review, and Update on Management. *Neurosurgical Focus* 2011;31(6):E15. DOI: 10.3171/2011.9.FOCUS11198.
85. Schaffler N, et al. Accuracy of Diagnostic Tests in Multiple Sclerosis--A Systematic Review. *Acta Neurologica Scandinavica* 2011;124(3):151-64. DOI: 10.1111/j.1600-0404.2010.01454.x.
86. Schoenfeld AJ, Bono CM, McGuire KJ, Warholc N, Harris MB. Computed Tomography Alone Versus Computed Tomography and Magnetic Resonance Imaging in the Identification of Occult Injuries to the Cervical Spine: A Meta-Analysis. *Journal of Trauma* 2010;68(1):109-13; discussion 113-4. DOI: 10.1097/TA.0b013e3181c0b67a.
87. Sellner J, et al. Diagnostic Workup of Patients with Acute Transverse Myelitis: Spectrum of Clinical Presentation, Neuroimaging and Laboratory Findings. *Spinal Cord* 2009;47(4):312-7. DOI: 10.1038/sc.2008.143.
88. Sicotte NL. Magnetic Resonance Imaging in Multiple Sclerosis: The Role of Conventional Imaging. *Neurologic Clinics* 2011;29(2):343-56. DOI: 10.1016/j.ncl.2011.01.005.
89. Sixta S, et al. Screening for Thoracolumbar Spinal Injuries in Blunt Trauma: An Eastern Association for the Surgery of Trauma Practice Management Guideline. *Journal of Trauma and Acute Care Surgery* 2012;73(5 Suppl 4):S326-32. DOI: 10.1097/TA.0b013e31827559b8.
90. Smith AB, Smirniotopoulos JG. Imaging Evaluation of Demyelinating Processes of the Central Nervous System. *Postgraduate Medical Journal* 2010;86(1014):218-29. DOI: 10.1136/pgmj.2009.087452.
91. Smith AB, Soderlund KA, Rushing EJ, Smirniotopolous JG. Radiologic-Pathologic Correlation of Pediatric and Adolescent Spinal Neoplasms: Part 1, Intramedullary Spinal Neoplasms. *American Journal of Roentgenology* 2012;198(1):34-43. DOI: 10.2214/AJR.10.7311.
92. Soderlund KA, Smith AB, Rushing EJ, Smirniotopolous JG. Radiologic-Pathologic Correlation of Pediatric and Adolescent Spinal Neoplasms: Part 2, Intradural Extramedullary Spinal Neoplasms. *American Journal of Roentgenology* 2012;198(1):44-51. DOI: 10.2214/AJR.11.7121. .
93. Sorantin E, Robl T, Lindbichler F, Riccabona M. MRI of the Neonatal and Paediatric Spine and Spinal Canal. *European Journal of Radiology* 2008;68(2):227-34. DOI: 10.1016/j.ejrad.2008.06.032.
94. Sucato DJ. Management of Severe Spinal Deformity: Scoliosis and Kyphosis. *Spine* 2010;35(25):2186-92. DOI: 10.1097/BRS.0b013e3181feab19.
95. Suspected Spine Trauma. ACR Appropriateness Criteria [Internet] American College of Radiology (ACR). 2012 Accessed at: <http://www.acr.org/>. [created 1999; accessed 2014 Sept 23]
96. Tekes A, Pinto PS, Huisman TA. Birth-Related Injury to the Head and Cervical Spine in Neonates. *Magnetic Resonance Imaging Clinics of North America* 2011;19(4):777-90; viii. DOI: 10.1016/j.mric.2011.08.004.
97. Thakkar RS, Malloy JP, Thakkar SC, Carrino JA, Khanna AJ. Imaging the Postoperative Spine. *Radiologic Clinics of North America* 2012;50(4):731-47. DOI: 10.1016/j.rcl.2012.04.006.
98. Thawait GK, Chhabra A, Carrino JA. Spine Segmentation and Enumeration and Normal Variants. *Radiologic Clinics of North America* 2012;50(4):587-98. DOI: 10.1016/j.rcl.2012.04.003.
99. Theodoridou A, Settas L. Demyelination in Rheumatic Diseases. *Postgraduate Medical Journal* 2008;84(989):127-32. DOI: 10.1136/jnnp.2005.075861.
100. Thwaites G, et al. British Infection Society Guidelines for the Diagnosis and Treatment of Tuberculosis of the Central Nervous System in Adults and Children. *Journal of Infection* 2009;59(3):167-87. DOI: 10.1016/j.jinf.2009.06.011.
101. Tilt L, Babineau J, Fenster D, Ahmad F, Roskind CG. Blunt Cervical Spine Injury in Children. *Current Opinion in Pediatrics* 2012;24(3):301-6. DOI: 10.1097/MOP.0b013e3283527035.
102. Toiviainen-Salo S, Markula-Patjas K, Kerttula L, Soini I, Valta H, Makitie O. The Thoracic and Lumbar Spine in Severe Juvenile Idiopathic Arthritis: Magnetic Resonance Imaging Analysis in 50 Children. *Journal of Pediatrics* 2012;160(1):140-6. DOI: 10.1016/j.jpeds.2011.06.030.
103. Tompkins M, Panuncialman I, Lucas P, Palumbo M. Spinal Epidural Abscess. *Journal of Emergency Medicine* 2010;39(3):384-90. DOI: 10.1016/j.jemermed.2009.11.001.
104. Tourdias T, et al. Assessment of Disease Activity in Multiple Sclerosis Phenotypes with Combined Gadolinium- and Superparamagnetic Iron Oxide-Enhanced MR Imaging. *Radiology* 2012;264(1):225-33. DOI: 10.1148/radiol.12111416.
105. Tracy JA, Bartleson JD. Cervical Spondylotic Myelopathy. *Neurologist* 2010;16(3):176-87. DOI: 10.1097/NRL.0b013e3181da3a29.

106. Trigylidas T, Yuh SJ, Vassilyadi M, Matzinger MA, Mikrogianakis A. Spinal Cord Injuries Without Radiographic Abnormality at Two Pediatric Trauma Centers in Ontario. *Pediatric Neurosurgery* 2010;46(4):283-9. DOI: 10.1159/000320134.
107. Tsujio T, et al. Characteristic Radiographic or Magnetic Resonance Images of Fresh Osteoporotic Vertebral Fractures Predicting Potential Risk for Nonunion: A Prospective Multicenter Study. *Spine* 2011;36(15):1229-35. DOI: 10.1097/BRS.0b013e3181f29e8d.
108. Van Tubergen A, Weber U. Diagnosis and Classification in Spondyloarthritis: Identifying a Chameleon. *Nature Reviews. Rheumatology* 2012;8(5):253-61. DOI: 10.1038/nrrheum.2012.33.
109. Vanhoenacker FM, Eyselbergs M, Cotten A. Spinal Degeneration: Beyond Degenerative Disc Disease: How to Discriminate Degeneration from Spondylarthropathies? *Neuroradiology* 2011;53 Suppl 1:S175-9. DOI: 10.1007/s00234-011-0936-1.
110. Wald JT. Imaging of Spine Neoplasm. *Radiologic Clinics of North America* 2012;50(4):749-76. DOI: 10.1016/j.rcl.2012.04.002.
111. Walker RC, Brown TL, Jones-Jackson LB, De Blanche L, Bartel T. Imaging of Multiple Myeloma and Related Plasma Cell Dyscrasias. *Journal of Nuclear Medicine* 2012;53(7):1091-101. DOI: 10.2967/jnumed.111.098830.
112. Wassenaar M, et al. Magnetic Resonance Imaging for Diagnosing Lumbar Spinal Pathology in Adult Patients with Low Back Pain or Sciatica: A Diagnostic Systematic Review. *European Spine Journal* 2012;21(2):220-7. DOI: 10.1007/s00586-011-2019-8.
113. Weber U, Lambert RG, Ostergaard M, Hodler J, Pedersen SJ, Maksymowych WP. The Diagnostic Utility of Magnetic Resonance Imaging in Spondylarthritis: An International Multicenter Evaluation of One Hundred Eighty-Seven Subjects. *Arthritis and Rheumatism* 2010;62(10):3048-58. DOI: 10.1002/art.27571.
114. Weber U, Maksymowych WP. Sensitivity and Specificity of Magnetic Resonance Imaging for Axial Spondyloarthritis. *American Journal of the Medical Sciences* 2011;341(4):272-7. DOI: 10.1097/MAJ.0b013e31820f8c59.
115. Weber U, Ostergaard M, Lambert RG, Maksymowych WP. The Impact of MRI on the Clinical Management of Inflammatory Arthritides. *Skeletal Radiology* 2011;40(9):1153-73. DOI: 10.1007/s00256-011-1204-5.